



Getting
started with

WAINUA®

Connect360®
Patient Support Program

See inside for important
information on the **Connect360**
Patient Support Program.



Table of Contents

Understanding hATTR PN (hereditary transthyretin amyloidosis associated polyneuropathy).....	4
Understanding WAINUA.....	6
My WAINUA Treatment	7
Getting Started with WAINUA.....	8
Possible Side Effects.....	10
Frequently Asked Questions	11

This brochure provides a summary of information about WAINUA, but it doesn't tell you everything.

If you need more information at any time, don't hesitate to ask your healthcare team. You can also reach out to the Connect360 Patient Support Program using the contact information on the **back of this brochure**.

Understanding hATTR PN

Hereditary transthyretin amyloidosis (or “hATTR”) is a rare, progressive condition

hATTR is caused by a “mutation” (change) in the gene that makes a protein in the body called “**transthyretin**” (or “**TTR**”).



Transthyretin (TTR)

hATTR happens when TTR proteins clump together and build up in the body to form abnormal structures called **amyloid fibrils**. The buildup of TTR can **stop the organs in your body from working properly**.



Amyloid fibrils

Sometimes this can lead to **damage to the body’s peripheral nerves**, a condition called “**polyneuropathy**” (PN).



Polyneuropathy

hATTR PN = hereditary transthyretin amyloidosis associated polyneuropathy

More information and resources on hATTR can be found at:

Hereditary Amyloidosis Canada

www.madhattr.ca

The Canadian Amyloidosis Support Network

www.amyloidosiscanada.org




Understanding WAINUA

WAINUA is used to treat polyneuropathy in adults who have stage 1 or stage 2 hATTR.



How does WAINUA work?

WAINUA belongs to a group of medicines called “antisense oligonucleotides” (ASO). It reduces the levels of TTR proteins in the body by preventing TTR from being made by the liver.



Learn how to take WAINUA

Scan the code to watch a video with step-by-step instructions on how to inject WAINUA



My WAINUA Treatment



How do I take WAINUA?

WAINUA is given as a subcutaneous injection (i.e., under the skin). It is packaged as a single-dose, pre-filled autoinjector. Each autoinjector can only be used one time.



How often do I take WAINUA?

The recommended dose of WAINUA is 45 mg (the contents of one autoinjector), once every month.

Always use WAINUA exactly as your healthcare professional has instructed you. Check with your healthcare professional if you are unsure. You and your healthcare professional will decide if you can inject WAINUA by yourself or if a caregiver should give you your injection. You and/or your caregiver will receive training on the right way to prepare and inject WAINUA.



Getting Started with **WAINUA**

It's common to have questions when starting treatment with WAINUA, such as:

- When can I get started?
 - What are my coverage options?
 - Who can I call for help?
-

This is where your **Connect360 Patient Support Program Care Coordinator** comes in.

Connect360
Patient Support Program 



Your Program Care Coordinator will complete your enrolment, answer any questions you may have, and help you get started on treatment.



Your first call

You will receive a welcome call from your Connect360 Patient Support Program Care Coordinator shortly after your physician has prescribed WAINUA and enrolled you in the program.

In preparation for this call, please have the following ready:

- Your health card
- Your health insurance benefits card and/or plan information, including the company name and policy number
- An overview of your medical history
- A list of current medications

Note: The caller ID should appear as “Connect360” so make sure to answer the phone if you see this name.

After your call, your Connect360 Patient Support Program Care Coordinator will process your information to complete your enrolment.



Your first injection

Once your enrolment is complete, the Connect360 Patient Support Program will schedule an appointment to provide training for your first WAINUA injection. During this appointment, you and your partner/caregiver will receive comprehensive training and support on how to inject WAINUA so that you can confidently receive injections at home moving forward.

Possible Side Effects

What are possible side effects from using WAINUA?

- Vomiting
- Nausea
- Blurred vision
- Injection site reactions (redness, itching, pain)
- Muscle aches or pain
- Rash
- Hair loss

These are not all the possible side effects that you may have when taking WAINUA. If you experience any side effects not listed here, tell your healthcare team.

What should I know about WAINUA and my vitamin A levels?

Treatment with WAINUA is expected to decrease your vitamin A levels. You will probably need to take a vitamin A supplement. Your healthcare team will monitor your vitamin A levels and tell you how much vitamin A you need to take. If you experience any symptoms of vitamin A deficiency, tell your healthcare team. Symptoms may include:

- Dry eyes
- Poor vision
- Decreased night vision
- Hazy or cloudy vision

Frequently Asked Questions

Where do I inject WAINUA?

You can inject WAINUA into the lower part of the stomach (abdomen) or the upper thighs, just under the skin. If a caregiver is giving you the injection, they may also inject WAINUA in the back of your upper arm.

What should I do if I miss a dose?

If you miss your dose of WAINUA, take your next dose as soon as possible. The dosing schedule for your monthly injections will continue from this date. If you have questions about your schedule, ask your healthcare team. Do not stop treatment with WAINUA unless your healthcare team tells you to.

How should I store WAINUA?

Store WAINUA in the refrigerator (2°C to 8°C) in the original carton to protect from light. Do NOT freeze or expose to heat.

If needed, WAINUA can be stored at room temperature (up to 30°C) in the original carton for up to 6 weeks. If it is not used within 6 weeks of storing at room temperature, discard the medication.

Keep out of reach and sight of children.

Can I take WAINUA if I am pregnant?

No. Tell your healthcare team right away if you become pregnant while taking WAINUA as you should stop your treatment. WAINUA affects your vitamin A levels, which can harm your unborn baby. If you are of childbearing potential, use an effective method of birth control while taking WAINUA.

The Connect360 Patient Support Program provides you with personalized resources and services throughout your treatment with WAINUA.



Dedicated Program Care Coordinator

- A dedicated Program Care Coordinator as a primary point of contact



Injection training and services

- Comprehensive, in-person training on how to inject WAINUA
- Monthly delivery of WAINUA to your home
- Regular reminders to help stay on track with treatment



Reimbursement navigation and financial assistance

- Exploration of your insurance and coverage options
- Assistance with minimizing out-of-pocket costs for your medication



Ongoing support and resources

- A Program Care Coordinator to help answer any questions about WAINUA
- Regular check-in calls throughout your treatment
- Access to a toll-free support line

If you have any questions about treatment with WAINUA, talk to your healthcare team or contact the Connect360 Patient Support Program at **1-833-733-ATTR (2887)** (Mon–Fri, 8 am–8 pm EST) or **support@connect360attr.ca**